

Synthesis of Donor-Acceptor copolymer By Using Benzoselenadiazole as acceptor for OTFT

Baji Shaik,^{a,1} Jin-Hee Han,^a Dong Jin Song,^a Hun-Min Kang,^a Ye Beyeol Kim,^{b,1} Chan Eon Park,^{b,*} Sang-Gyeong Lee^{a,*}

^aDepartment of Chemistry, Research Institute of Natural Science (RINS), Graduate School for Molecular Materials and Nanochemistry, Gyeongsang National University, Jinju 660-701, Korea.

^bDepartment of Chemical Engineering Pohang University of Science and Technology POSTECH Organic Electronics Laboratory Polymer Research Institute, Pohang, 790-784 (Korea)

*Corresponding authors:

Sang-Gyeong Lee: Tel: +82 55 772 1487; fax: +82 55 772 1489; E-mail address: leesang@gnu.ac.kr

Chan Eon Park: E-mail: cep@postech.ac.kr

Table S1. 2D-GIXD characteristic properties.

	$q[\text{\AA}^{-1}]$ (d-spacing $[\text{\AA}]$)	
	(100)	π - π
ascast	0.261 (24.1)	1.72 (3.65)
Ann 120	0.256(24.5)	1.72 (3.65)
Ann 150	0.276(22.8)	1.72 (3.65)
Ann 180	0.281(22.4)	1.71(3.67)