

Supplementary Information

Synergistic assembly of nanoparticles in smectic liquid crystals

Apiradee Honglawan,^[a] Dae Seok Kim^[b], Daniel Beller,^{[c],†} Dong Ki Yoon^[b], Mohamed A. Gharbi,^[a,c, d] Kathleen J. Stebe,^[a] Randall D. Kamien^[c] and Shu Yang ^{* [a,d]}

^aDepartment of Chemical and Biomolecular Engineering, University of Pennsylvania, Towne Building, 220 South 33rd Street, Philadelphia, PA 19104, USA

^b Graduate School of Nanoscience and Technology, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 305-701, Republic of Korea

^cDepartment of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA 19104, USA

^dDepartment of Materials Science and Engineering, 3231 Walnut Street, University of Pennsylvania, Philadelphia, PA 19104, USA

[†]Current address: School of Engineering and Applied Sciences, Harvard University, 29 Oxford Street, Cambridge, MA 02138, USA

*Corresponding author: Shu Yang (shuyang@seas.upenn.edu), 215-898-9645

Video S1. *In situ* observation of the mixture of F-SiO₂ NP (100 nm)/F-LC cooled from isotropic phase (200 °C) to SmA phase (114 °C) under polarized optical microscope (POM), revealing a gradual transformation from an array of TFCDs to intermediate states of mixed TFCDs and fan textures to eventually fan textures.