Supporting information for:

## Nematic field transfer in a two-dimensional protein fibril assembly

Sophia Jordens,<sup>1</sup> Konrad Schwenke,<sup>2</sup> Ivan Usov,<sup>1</sup> Emanuela Del Gado,<sup>3,2,\*</sup> and Raffaele Mezzenga<sup>1,\*</sup>

<sup>1</sup>Department of Health Sciences & Technology, ETH Zurich, Schmelzbergstrasse 9, 8092 Zurich, Switzerland

<sup>2</sup>Department of Civil, Environmental & Geomatic Engineering, ETH Zurich, Stefano-

Franscini-Platz 3, 8093 Zurich, Switzerland

<sup>3</sup>Institute for Soft Matter Synthesis and Metrology, Georgetown University, 37<sup>th</sup> and O

Streets, N.W., Washington D.C. 20057, USA

\*ed610@georgetown.edu

\*raffaele.mezzenga@hest.ethz.ch



Figure S1. AFM image of fibrils and short aggregates adsorbed at the air-liquid interface of a solution containing both species at concentrations  $c_f = 0.01\%$  w/w and  $c_a = 0.001\%$  w/w.



Figure S2. AFM image of fibrils and short aggregates adsorbed at the air-liquid interface of a solution containing both species at concentrations  $c_f = 0.005\%$  w/w and  $c_a = 0.01\%$  w/w.



Figure S3. AFM image of short aggregates adsorbed at the air-liquid interface of a solution containing both species at concentrations  $c_a = 0.01\%$  w/w.