

Supporting Information

Magneto-Fluorescent Nanoparticles with High-Intensity NIR Emission, T_1 and T_2 Weighted MR for Multimodal Specific Tumor Imaging

Guannan Wang,^{ac#} Xuanjun Zhang,^{b#} Yaxu Liu,^a Zhangjun Hu,^c Xifan Mei,^{a*} Kajsa Uvdal^{*c}

#These two authors contributed equally to this work.

^aCollege of Pharmacy & the Key Laboratory for Medical Tissue Engineering of Liaoning Province, Liaoning Medical University, Jinzhou, 121001, China. E-mail: meixifan1971@163.com; Fax: +86-416-4673162; Tel: +86-416-4673162

^bFaculty of Health Sciences, University of Macau, Avenida da Universidade, Taipa, Macau, China

^cDivision of Molecular Surface Physics & Nanoscience, Department of Physics, Chemistry, and Biology, Linköping University, Linköping 58183, Sweden. E-mail: kajsa@ifm.liu.se; Fax: +46 13 28 8969; Tel: +46 13281208




Figure S1. The TGA of FMNPs




Figure S2. Flow cytometry histograms of pure A549 cancer cells (Blue) and the cells after one hour incubation with the FMNPs (green) suspension at 0.5 mg/mL.