Substituent groups effect on the morphology and memory performance of phenazine derivatives

Pei-Yang Gu,¹ Yong Ma,² Jing-Hui He,² Guankui Long,¹ Chengyuan Wang,¹ Wangqiao Chen,¹ Yi Liu,³ Qing-Feng Xu,^{2, *} Jian-Mei Lu,^{2, *} and Qichun

Zhang^{1,4*}

¹ School of Materials Science and Engineering, Nanyang Technological University, Singapore 639798, Singapore. E-mail: gczhang@ntu.edu.sg

²College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou, 215123, China. E-mail: lujm@suda.edu.cn;

xuqingfeng@suda.edu.cn

³The Molecular Foundry and [‡]Advanced Light Source, Lawrence Berkeley National Laboratory, One Cyclotron Road, Berkeley, California 94720, United States ⁴Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

Contents

Figure S1 ¹H NMR spectrum of 2OHPz in CF₃COOD (TFA).

Figure S2 ¹³C NMR spectrum of 2OHPz in TFA.

Figure S3 HRMS spectrum of 20HPz.

Figure S4¹H NMR spectrum of 10HPz in CDCl₃.

Figure S5¹³C NMR spectrum of 10HPz in CDCl₃.

Figure S6 HRMS spectrum of 10HPz.

Figure S7 TGA curves of 20HPz and 10HPz with a heating rate of 20 °C min⁻¹ under

N₂.

Figure S8 Stability of the memory device (ITO/2OHPz/Al and ITO/1OHPz/Al,

respectively) under a constant "read" voltage of -1V.

Figure S9 Normalized optical absorption spectra of 2OHPz and 1OHPz thin films on

a quartz plate.

Figure S10 Cyclic voltammogram of the ferrocene standard in dichloromethane solution at the scan rate of 100 mV S⁻¹.



Figure S2 ¹³C NMR spectrum of 2OHPz in TFA.



Figure S4 ¹H NMR spectrum of 10HPz in CDCl₃.



Figure S6 HRMS spectrum of 10HPz.



Figure S7 TGA curves of 20HPz and 10HPz with a heating rate of 20 °C min⁻¹ under



Figure S8 Stability of the memory device (ITO/2OHPz/A1 and ITO/1OHPz/A1, respectively) under a constant "read" voltage of -1V.



Figure S9 Normalized optical absorption spectra of 2OHPz and 1OHPz thin films on a quartz plate.



Figure S10 Cyclic voltammogram of the ferrocene standard in dichloromethane solution at the scan rate of 100 mV S^{-1} .