

Green-blue light-emitting platinum(II) complexes of cyclometallated 4,6-difluoro-1,3-dipyridylbenzenes showing mesophase organisation

**Valery N. Kozhevnikov^{[a]*}, Bertrand Donnio^{[b,c]*},
Benoît Heinrich^[b], J. A. Gareth Williams^[d] and Duncan W. Bruce^{[e]*}**

^a *School of Life Sciences, Northumbria University, Newcastle-Upon-Tyne, NE1 8ST, UK. Email: valery.kozhevnikov@northumbria.ac.uk*

^b *Institut de Physique et Chimie des Matériaux de Strasbourg (IPCMS), UMR 7504, CNRS-Université de Strasbourg, 23 rue du Loess, BP43, 67034 Strasbourg cedex 2, France. Email: bertrand.donnio@ipcms.unistra.fr*

^c *Complex Assemblies of Soft Matter Laboratory (COMPASS), UMI 3254, CNRS-Solvay-University of Pennsylvania, CRTB, 350 George Patterson Boulevard, Bristol, PA 19007, USA*

^d *Department of Chemistry, Durham University, Durham. DH1 3LE, UK.*

^e *Department of Chemistry, University of York, Heslington, YORK. YO10 5DD UK. E-mail: duncan.bruce@york.ac.uk*

Supplementary Information

DSC data




Figure S1: DSC of **7** (1.51 mg, rate 2°C.min⁻¹): 1st heating (top), 1st cooling (bottom).




Figure S2: DSC of **5** (1.00 mg, rate 5°C.min⁻¹): 1st heating (top), 1st cooling (bottom).




Figure S3: SAXS diffractogram of **7** in the three-dimensional mesophase at 60°C.

Luminescence spectra in dichloromethane solution.

