

Supporting Information

Calix[4]arene crownether as an oriented linker for highly sensitive detection of zinc ions using peptide probe

Shengsong Jia^{a,b}, Min Shao^c, Fei Zou^b, Beiping Wu^d, Hongjian Zhou^c, Hongxia Chen^{a,b,*}

- State Key Laboratory of Dairy Biotechnology, Bright Dairy & Food Co. Ltd., Shanghai 200444, China
- Laboratory of Biosensing Technology, School of Life Sciences, Shanghai University, Shanghai 200444, China
- Instrumental Analysis and Research Center, School of Materials Science and Engineering, Shanghai University, Shanghai 200444, China
- Shanghai Key Laboratory of Bio-Energy Crops, School of Life Sciences, Shanghai University, Shanghai 200444, China
- Key Laboratory of Materials Physics, Centre for Environmental and Energy Nanomaterials, Anhui Key Laboratory of Nanomaterials and Nanotechnology, Institute of Solid State Physics, Chinese Academy of Sciences, Hefei 230031, China




Figure S1 The original SPR data for real samples detection. (a) Urine, (b) Tap water and (c) Pond water: black line (0.1 μM) and red line (1 μM).

*Corresponding author: Tel.: +86 21 66137539. E-mail address: hxchen@shu.edu.cn (H.Chen)