

Table S2.Stability, intra-day and inter-day precision of the proposed method

Analyte	Stability (n = 6)		Intra-day (n=6)		Inter-day (n=6)	
	Mean ± SD	RSD (%)	Mean ± SD	RSD (%)	Mean ± SD	RSD (%)
	µg/mL		µg/mL		µg/mL	
3'-Hydroxypuerarin	136.82±0.99	0.73	140.26±1.37	0.98	137.66±1.04	0.76
Puerarin	918.34±8.48	0.92	924.26±5.91	0.64	932.00±11.10	1.19
3'-Methoxypuerarin	637.36±4.09	0.64	638.87±4.81	0.75	639.23±7.28	1.14
Daidzin	87.87±0.91	1.04	87.78±0.88	1.01	88.35±0.99	1.12
Rutin	48.17±0.51	1.06	48.35±0.25	0.52	48.34±0.44	0.92
Astragalin	13.62±0.10	0.74	13.58±0.05	0.34	13.62±0.07	0.55
Daidzein	19.48±0.34	1.77	18.16±0.30	1.64	17.85±0.21	1.16
Ginsenoside Rg ₁	160.03±1.22	0.77	163.02±2.37	1.45	161.18±1.15	0.71
Astragaloside IV	311.47±3.88	1.24	305.99±1.30	0.42	307.06±2.70	0.88
20(S)-Ginsenoside Rg ₃	44.56±0.54	1.22	45.58±0.45	0.98	44.93±0.45	1.00