

Supporting Information

The two datasets used in this article were listed:

The first metabolomics data contained three classes, healthy control C57 mice, female C57-AMPK gene knocked-out mice and male C57-AMPK gene knocked-out mice. The concentrations of 42 compounds were listed in Table 1.

Table 1. Qualitative and quantitative metabolic profile of three group mice.

id	t _r ^a (min)	endogenous metabolites	C57	AMPK-male	AMPK-female
1	5.922	Aminoethane	0.2456±0.0705	0.1905±0.0567	0.1958±0.0551
2	6.593	Ethylene glycol	0.0182±0.0020	0.0530±0.0428	0.0746±0.0626
3	6.84	N,N-diethylacetamide	0.0657±0.0087	0.0476±0.0202	0.0557±0.0107
4	7.716	Lactic acid *	0.0872±0.0374	0.0952±0.0592	0.1482±0.2155
5	7.934	Acetic acid	0.0856±0.0333	0.0229±0.0140	0.0412±0.0203
6	10.01	phosphate	2.1278±0.9173	1.4730±0.7381	1.3767±0.9361
7	10.2	l-Threonine	0.0173±0.0098	0.0108±0.0068	0.0096±0.0065
8	10.297	Phenylacetic acid	0.0047±0.0023	0.0159±0.0103	0.0147±0.0097
9	10.382	Succinic acid *	0.0311±0.0129	0.0098±0.0031	0.0119±0.0086
10	10.447	1,2-Hydroquinone	0.0120±0.0072	0.0078±0.0047	0.0067±0.0039
11	10.503	Glyceric acid	0.0961±0.0266	0.0400±0.0232	0.0183±0.0087
12	10.723	(R*,R*)-2,3-Dihydroxybutano	0.0167±0.0053	0.0037±0.0014	0.0053±0.0029

		ic acid			
13	11.357	2,4-Dihydroxybutanoic acid	0.0147±0.0051	0.0155±0.0080	0.0166±0.0047
14	11.583	(R*,S*)-3,4-Dihydroxybutanoic acid	0.0304±0.0098	0.0132±0.0064	0.0178±0.0107
15	11.797	N-(1-oxobutyl)- Glycine	0.0653±0.0244	0.0319±0.0186	0.0274±0.0151
16	12.341	Isovaleroglycine	0.0356±0.0134	0.0160±0.0079	0.0107±0.0073
17	12.483	D- Threitol	0.0714±0.0273	0.0290±0.0130	0.0251±0.0151
18	12.645	N-Crotonylglycine	0.0240±0.0146	0.0207±0.0129	0.0148±0.0099
19	12.973, 13.203	2,3,4- Trihydroxybutyrate	0.1276±0.0162	0.0631±0.0343	0.0412±0.0250
20	14.53	N-(1-oxohexyl)-glycine	0.0960±0.0319	0.0421±0.0273	0.0232±0.0081
21	14.58	3-Hydroxyphenylacetic acid	0.0326±0.0100	0.0140±0.0081	0.0134±0.0088
22	14.713	D-Xylose	0.0408±0.0150	0.0182±0.0044	0.0193±0.0053
23	14.823, 15.057	D-Ribose	0.0926±0.0370	0.0252±0.0142	0.0250±0.0179
24	15.509, 15.733	Arabitol	0.0287±0.0164	0.0283±0.0179	0.0278±0.0215
25	16.023	6-Deoxy-D-Galactose,	0.0336±0.0083	0.0177±0.0100	0.0149±0.0104
26	16.087	Mannonic acid	0.0505±0.0177	0.0211±0.0143	0.0168±0.0138
27	16.2	cis-Aconitic acid*	0.0535±0.0288	0.0105±0.0079	0.0168±0.0147
28	16.357	Phosphoric acid	0.0414±0.0202	0.0230±0.0141	0.0212±0.0168
29	17.177	Isocitric acid*	0.0348±0.0121	0.0140±0.0093	0.0248±0.0138
30	17.563	Hippuric acid	0.0470±0.0126	0.0180±0.0074	0.0156±0.0096
31	17.85, 17.96	D-Fructose*	0.0512±0.0286	0.0371±0.0145	0.0480±0.0131
32	18.087	N-phenyl glycine*	0.0596±0.0214	0.0455±0.0272	0.0389±0.0287

33	18.197, 18.147	D-Glucose*	0.3785±0.1618	0.1741±0.0654	0.1859±0.0736
34	18.507	Altronic acid	0.0302±0.0069	0.0185±0.0100	0.0102±0.0074
35	18.577, 18.65	D-Sorbitol*	0.0896±0.0269	0.0254±0.0187	0.0300±0.0275
36	18.983, 19.533	Galactonic acid	0.0613±0.0282	0.0617±0.0328	0.0441±0.0351
37	19.99	Palmitic acid	0.0084±0.0009	0.0067±0.0017	0.0071±0.0025
38	20.403	Myo-Inositol	0.0347±0.0228	0.0097±0.0037	0.0134±0.0129
39	25.465	D-Turanose	0.0216±0.0138	0.0197±0.0090	0.0510±0.0099
40	25.653, 25.783	D- (+) -lactose monohydrate*	1.0400±0.3349	0.7475±0.2366	0.6559±0.3286
41	25.927	Lactose*	0.0142±0.0043	0.0143±0.0075	0.0190±0.0163

*: Identified by standard substances; ^a: Retention time;

The second data contained 25 mice. i.e., 20 KK-Ay mice with diabetes and 5 C57 mice as healthy control group. 20 KK-Ay diabetic mice were further classified into four classes, 5 mice in each class. Each group was treated with repaglinide with different periods ranging from 14th week to 22nd week. 42 compounds were considered as endogenous metabolites to represent each sample. The concentrations of 42 compounds were listed in Table 2.

Table 2 Quantitative metabolic profile of the repaglinide treated diabetic KK-A^y mice and C57BL/6J mice (relative quantity of each metabolite to internal standard).

id	t _r ^a (min)	endogenous metabolites	14th week	17th week	20th week	22nd week	healthy
1	6.833	N,N-diethyl-acetamide	6.59±0.70	6.85±1.14	7.54±0.48	8.06±0.45	5.19±0.84
2	7.14	N-ethyl-acetamide	15.42±2.70	14.72±2.93	16.32±0.69	20.17±2.03	11.90±1.42
3	7.667	Lactic acid	30.09±11.02	165.99±46.18	46.61±26.27	66.03±11.22	6.25±3.97
4	7.75	Dimethylsilanediol	6.59±1.16	6.67±1.02	5.70±2.17	6.63±3.15	3.83±0.68
5	7.883	Glycollic acid	3.24±0.86	3.03±1.73	4.07±1.39	4.09±1.03	7.45±1.62

6	8.557	N-formyl-glycine	6.10±0.79	12.99±2.01	16.65±3.09	9.59±7.84	9.81±1.62
7	8.993	Valine	8.46±5.08	15.88±2.27	15.26±2.62	6.78±7.25	7.49±0.44
8	10.32	Butanedioic acid	6.58±7.26	4.04±0.75	3.33±0.18	5.80±3.12	7.70±3.62
9	10.37	o-Dihydroxybenzene	1.03±0.26	0.64±0.22	0.78±0.48	1.38±0.92	0.51±0.18
10	10.427	2,3-Bis(hydroxy)propionic acid	1.13±0.16	0.93±0.53	0.99±0.41	1.44±0.25	9.26±2.15
11	11.247	Pentanedioic acid	1.83±0.29	1.98±0.30	4.60±2.82	2.37±0.58	0.79±0.30
12	12.163	Malic acid	2.80±1.11	1.46±0.41	1.49±0.86	1.78±0.47	0.94±0.51
13	12.37	Threitol	3.87±0.47	2.84±0.90	4.58±0.84	4.01±1.01	6.77±0.46
14	12.857	2,3,4-Trihydroxybutyric acid*	2.15±0.85	1.41±0.34	1.93±0.31	1.73±0.77	5.74±0.74
15	13.083	2,3,4-Trihydroxybutyric acid*	1.41±0.41	1.42±0.69	2.00±0.71	1.67±0.55	6.20±0.97
16	13.373	α-Hydroxyglutaric acid	3.93±1.99	5.01±1.13	3.38±1.13	2.94±1.89	6.91±6.44
17	14.523	2-Phenyl-1,2-bis(hydroxy)propane	1.58±0.25	1.91±2.85	0.79±0.70	2.00±1.08	0.42±0.21
18	15.377	Arabitol	1.99±0.24	1.47±0.67	2.86±1.39	2.11±0.57	2.36±0.47

19	15.597	Xylitol	1.17±0.18	1.45±1.18	1.62±0.59	1.80±0.71	2.37±0.36
20	15.893	2,3-Butanediol	1.68±0.56	1.08±0.58	0.34±0.17	1.42±0.70	0±0 ^d
21	15.967	Pentonic acid	1.52±0.43	0.54±0.34	1.17±1.15	1.72±1.05	3.56±0.20
22	16.1	1-Propene-1,2,3-tricarboxylic acid	3.79±1.19	2.15±1.36	3.44±2.00	2.32±1.16	6.95±1.02
23	17.493	1,5-Anhydro-d-glucitol ¹	42.31±5.74	15.12±11.53	5.20±3.45	0±0 ^b	3.48±0.88
24	17.73	d-Ribose	9.12±4.39	0.48±0.26	2.25±2.94	0.71±0.25	0.84±0.38
25	17.85	Xylitol	6.42±1.70	3.07±1.98	3.02±2.11	0.74±0.26	4.01±1.35
26	18.227	d-Glucose ²	1154.25±145.85	884.53±157.43	223.74±111.73	8.52±1.02	4.05±0.44
27	18.403	d-Galactose ³	379.15±54.26	354.68±128.86	68.94±41.86	1.64±0.21	2.76±0.14
28	18.56	d-Mannose	5.77±3.86	4.99±1.74	6.70±2.90	4.63±4.08	10.60±2.81
29	18.877	Ribonic acid	3.09±2.07	2.63±0.70	2.78±0.86	0.63±0.23	10.77±0.88
30	19.04	Glucopyranose	41.78±17.67	66.13±60.23	3.42±3.68	0.05±0.12	0±0 ^e
31	19.418	Tartaric acid	3.40±1.08	2.70±1.60	1.40±0.76	0.67±0.80	1.18±0.06

32	19.91	Hexadecanoic acid	2.18±0.83	2.77±1.10	3.73±2.03	0.53±0.91	0.39±0.17
33	20.277	Myo-inositol	4.39±1.37	3.46±1.27	2.88±0.87	2.65±1.29	1.79±0.40
34	21.584	9,12-Octadecadienoic acid	1.36±0.32	0.85±0.35	1.34±1.05	0±0 ^c	0±0 ^f
35	25.143	α-d-Glucopyranoside	49.0±23.9	9.69±9.12	21.52±23.40	19.73±21.06	5.52±3.65
36	25.54	d-Turanose [#]	56.9±12.9	63.02±8.29	56.97±5.63	39.15±17.90	85.91±26.03
37	25.667	Maltose	13.4±4.43	13.79±2.43	13.13±2.64	8.20±4.79	20.89±6.80
38	25.733	2-Deoxy-galactopyranose	2.13±0.85	0.60±0.37	0.57±0.35	4.02±6.57	1.29±0.43
39	25.903	d-Turanose [#]	25.7±6.98	8.03±7.20	11.51±8.56	9.62±11.23	1.98±1.10
40	26.04	Melibiose	3.07±0.89	4.61±2.70	2.07±1.05	2.11±1.44	0.19±0.06
41	26.107	Allonic acid	3.16±1.02	2.91±3.73	1.98±1.54	2.30±1.73	0.36±0.13

1,2,3: Identified by standard substances; a: Retention time;

b,c,d,e,f: No target product was detected

*,#:Can't identified by NIST05.