

## Support Information

### Monitoring Early Zeolite Formation via *in situ* Electrochemical Impedance Spectroscopy

G. Brabants,<sup>a,b</sup> S. Lieben,<sup>a</sup> E. Breynaert,<sup>a</sup> E. K. Reichel,<sup>b</sup> F. Taulelle,<sup>a</sup> J. A. Martens,<sup>a</sup> B. Jakoby,<sup>b</sup> and C. E. A. Kirschhock<sup>a</sup>

<sup>a</sup>Center for Surface Chemistry and Catalysis, KU Leuven, Leuven, Belgium.

<sup>b</sup>Institute for Microelectronics and Microsensors, Johannes Kepler University, Linz, Austria.

E-mail: [christine.kirschhock@biw.kuleuven.be](mailto:christine.kirschhock@biw.kuleuven.be)




Fig. 8. Small angle X-ray scattering data of the clear solutions with  $\text{Si}/\text{OH} = 4.64$  (upper) and  $\text{Si}/\text{OH} = 2.37$  (lower) fitted with four different models: Gaussian spheres, core/shell spheres, cylinders and the core/shell oblate ellipsoids proposed by Fedeyko et al.<sup>14</sup>. On the right side a detail is shown of the fitting in the higher  $q$  region: only the core/shell oblate ellipsoid model fits the experimental data well.