

Electronic Supplementary Information 2 (ESI)


Crystallisation, thermal analysis and acetal protection activity of new layered Zn(II) hybrid polymorphs

Salem Saïd,^a Houcine Naïli^{a*}, Thierry Bataille^b and Raquel P. Herrera^c

^a *Laboratoire Physico-chimie de l'Etat Solide, Département de Chimie, Faculté des Sciences de Sfax, B.P. 1171, 3000 Sfax, Université de Sfax, Tunisie. E-mail address: houcine_naili@yahoo.com*

^b *Laboratorio de Organocatalisis Asimétrica, Departamento de Química Orgánica, Instituto de Síntesis Química y Catalysis Homogénea (ISQCH), CSIC-Universidad de Zaragoza, C/Pedro Cerbuna 12, E-50009 Zaragoza, Spain.*

Figure S1. ^1H and ^{13}C -APT NMR spectra (CDCl_3) of compound **4a**.







Figure S2. ^1H and ^{13}C -APT NMR spectra (CDCl_3) of compound 4b.







Figure S3. ^1H and ^{13}C -APT NMR spectra (CD_2Cl_2) of compound **4c**.







Figure S4. ¹H and ¹³C-APT NMR spectra (CD₂Cl₂) of compound **4d**.






Figure S5. ^1H and ^{13}C -APT NMR spectra (CDCl_3) of compound **4e**.




Figure S6. ^1H and ^{13}C -APT NMR spectra (CDCl_3) of compound **4f**.




Figure S7. ^1H and ^{13}C -APT NMR spectra (CD_2Cl_2) of compound **4g**.




Figure S8. ^1H and ^{13}C -APT NMR spectra (CD_2Cl_2) of compound **4i**.

