

Electronic Supplementary Information

Phase Transitions in Hydrophobe/Phospholipid Mixtures: Hints at Connections between Pheromone and Anaesthetic Action

Silvia Borsacchi,^a Marco Geppi,^b Sara Macchi,^b Barry W. Ninham,^c Emiliano Fratini,^d Moira Ambrosi,^d Piero Baglionid,^e and Pierandrea Lo Nostro*^{d,e}

^a: Istituto di Chimica del Composti OrganoMetallici (ICCOM) del CNR, 56124 Pisa, Italy.

^b: Department of Chemistry and Industrial Chemistry, University of Pisa, 56124 Pisa, Italy.

^c: Department of Applied Mathematics, The Australian National University, Canberra ACT0200, Australia.

^d: Department of Chemistry & CSGI, University of Florence, 50019 Sesto Fiorentino (Firenze), Italy.

^e: Enzo Ferroni Foundation, 50019 Sesto Fiorentino (Firenze), Italy.




Figure S1. DSC runs for the different DOPC/pheromone samples as a function of the pheromone/DOPC mole ratio (c): 0 (●), 0.1 (■), 0.2 (▲), 0.7 (●), 0.9 (●), 1.8 (●), 7.3 (■). Curves have been offset for graphical purposes.