Electronic Supporting Information for:

## A 1,2,3-Dithiazolyl-*o*-naphthoquinone: A Neutral Radical with Isolable Cation and Anion Oxidation States

Chad S. Smithson,<sup>*a*</sup> Daniel J. MacDonald, <sup>*a*</sup> T. Matt Letvenuk,<sup>*a*</sup> Christian E. Carello,<sup>*a*</sup> Michael Jennings,<sup>*b*</sup> Alan Lough,<sup>*c*</sup> James Britten,<sup>*d*</sup> Andreas Decken,<sup>*e*</sup> and Kathryn E. Preuss\*<sup>*a*</sup>

<sup>a</sup>Department of Chemistry, University of Guelph, Guelph, Ontario, Canada N1G 2W1. <sup>b</sup>FreeLance Crystallography, 185 Chelsea Avenue, London, Ontario, Canada N6J 3J5. <sup>c</sup>Department of Chemistry, University of Toronto, Toronto, Ontario, Canada M5S 3H6. <sup>d</sup>Department of Chemistry, McMaster University, Hamilton, Ontario, Canada L8S 4M1. <sup>e</sup>Department of Chemistry, University of New Brunswick, Fredericton, NB, Canada E3B 5A3.



**Figure S1.** Experimental and simulated EPR spectra of neutral radical **1** in CH<sub>2</sub>Cl<sub>2</sub> at ambient temperature; simulation parameters:  $a_N = 2.934$  G,  $2 \times a_H = 0.386$  G,  $2 \times a_H = 0.299$  G, line width = 0.234 mT; correlation > 0.999.



**Figure S2.** CV of **1** in THF (0.1 M  $[nBu_4N][PF_6]$ ); multiple cycles with the anodic direction swept first; (top to bottom) progressively more negative reduction potentials in the cathodic direction



**Figure S3.** CV of **3** in CH<sub>3</sub>CN (0.10 M [*n*Bu<sub>4</sub>N][PF<sub>6</sub>]).