

Electronic Supporting Information

Polymer *versus* phosphine stabilized Rh nanoparticles as components of supported catalysts: implication in the hydrogenation of cyclohexene model molecule

M. Ibrahim,^{a,b} M. A. S. Garcia,^c L. L. R. Vono,^{a,b,c} M. Guerrero,^{a,b} P. Lecante,^d L. M. Rossi,^{*c} and K. Philippot^{*a,b}

^a CNRS, LCC (*Laboratoire de Chimie de Coordination*), 205 route de Narbonne, BP 44099, F-31077-Toulouse Cedex 4, France

^b Université de Toulouse, UPS, INPT, F-31077-Toulouse Cedex 4, France

^c Department of Fundamental Chemistry, Institute of Chemistry, University of São Paulo, PO Box 26077, 05513-970, São Paulo, SP, Brazil

^d CNRS UPR 8011; CEMES (Centre d'Elaboration des Matériaux et d'Etudes Structurales) 29 rue Jeanne Marvig, F-31055 Toulouse, France

* karine.philippot@lcc-toulouse.fr




Figure S1. TON *versus* Time curves obtained for Rhdppb, RhPPh₃ and RhPVP colloidal systems




Figure S2. TON *versus* Time curves obtained for RhPVP@FFSiNH₂ nanocatalyst up to 5 recycles




Figure S3. TON *versus* Time curves obtained for RhPPh₃@FFSiNH₂ nanocatalyst up to 5 recycles




Figure S4. TON *versus* Time curves obtained for Rhdppb@FFSiNH₂ nanocatalyst up to 5 recycles