

ELECTRONIC SUPPLEMENTARY MATERIAL OF THE ARTICLE:
**Cocoplum (*Chrysobalanus icaco* L.) anthocyanins exert anti-inflammatory
activity in human colon cancer and non-malignant colon cells**


Food & Function

**Vinicius P. Venancio^{1,2}, Paula A. Cipriano¹, Hyemee Kim¹, Lusânia M. G.
Antunes², Stephen T. Talcott¹, Susanne U. Mertens-Talcott^{1,*}**

¹ Department of Nutrition and Food Science, Texas A&M University, College
Station, Texas 77843-2254, USA

² Department of Clinical Analyses, Toxicology and Food Science, University of São
Paulo, Ribeirão Preto, São Paulo, 14040-903, Brazil

* Author to whom correspondence should be addressed [phone (979) 458-1819; fax
(979) 458-3704; e-mail: smtalcott@tamu.edu]



Supplementary Figure 1 – Anthocyanin chromatographic profile of CP extract. HPLC-ESI-MSⁿ analysis allowed the characterization of each peak as follows: delphinidin-3-glucoside (peak 1), cyanidin 3-glucoside (peak 2), petunidin 3-glucoside + delphinidin 3-(6''-acetyl)galactoside or delphinidin 3-(6''-oxaloyl)arabinoside (peak 3), peonidin 3-glucoside (peak 4), petunidin 3-(6''-acetyl)galactoside or petunidin 3-(6''-oxaloyl)arabinoside (peak 5) and peonidin 3-(6''-acetyl)glucoside or peonidin 3-(6''-oxaloyl)arabinoside (peak 6).