

Supporting Information

Can Self-Assembled Hydrogels Composed of Aromatic Amino Acid Derivative function as drug delivery carrier?

P. Tiwari,^a R. Rajagopalan,^b M. Moin,^b R. Soni,^b P. Trivedi^{b*} and A.Dutt Konar,^{a,b*}

^aDept of Chemistry, ^bSchool of Pharmaceutical Sciences , Rajiv Gandhi Technological University, Bhopal- 462033, MadhyaPradesh, India, E-mail: anitaduttkonar@rgtu.net, Ph:917554056278; E-mail: piyush.trivedi@rgtu.net Ph:917552742001

Contents:	Page Number
Camera view of Hydrogelator I and II	2
IR spectra of Hydrogelator I and II	3
¹H NMR of Hydrogelator I and II	4
Mass Spectra of Hydrogelator I and II	5




Figure S1. Camera view of the hydrogels obtained from a) Gelator **I** and b) **II**




Figure S2. IR Spectra of xerogel -I




Figure S2. IR Spectra of xerogel -II




Figure S3. ^1H NMR Spectra of Molecule -I




Figure S4. ^1H NMR Spectra of Molecule -II




Figure S5. Mass Spectra of Molecule-I




Figure S6. Mass Spectra of Molecule-II