

## Supporting Information

# Synthesis, and Optical and Electrochemical Properties of Water-Soluble Cationic Fluorophors Based on Bispyridinium and Dibenzothiophene-S,S-Dioxide

Guiting Chen,<sup>a</sup> Ruifeng He,<sup>a</sup> Wei Yang,<sup>a</sup> Bin Zhang\*<sup>ab</sup>

<sup>a</sup> *Institute of Polymer Optoelectronic Materials and Devices, State Key Laboratory of Luminescent Materials and Devices, South China University of Technology, Guangzhou 510640, China. E-mail: msbzhang@scut.edu.cn*

<sup>b</sup> *Key Laboratory of Optoelectronic Devices and Systems of Ministry of Education and Guangdong Province, College of Optoelectronic Engineering, Shenzhen University, Shenzhen 518060, China*




Figure S1. <sup>1</sup>H and <sup>13</sup>C NMR of **1**




Figure S2. <sup>1</sup>H and <sup>13</sup>C NMR of 2




Figure S3. <sup>1</sup>H and <sup>13</sup>C NMR of 3




Figure S4. <sup>1</sup>H and <sup>13</sup>C NMR of 4




Figure S5. <sup>1</sup>H and <sup>13</sup>C NMR of 5




Figure S6. <sup>1</sup>H and <sup>13</sup>C NMR of 6




Figure S7. <sup>1</sup>H and <sup>13</sup>C NMR of 8






Figure S8. <sup>1</sup>H and <sup>13</sup>C NMR of **9**





Figure S9. <sup>1</sup>H and <sup>13</sup>C NMR of FSOPyCl




Figure S10. <sup>1</sup>H and <sup>13</sup>C NMR of FSOmiCl