

Supporting Information

Facile synthesis of zwitterionic polyglycerol dendrimers with a β -cyclodextrin core as MRI contrast agent carriers

Yuxin Han^a, Yue Qian^b, Xiaoxuan Zhou^b, Hongjie Hu^b, Xiangrui Liu^a, Zhuxian Zhou^a, Jianbin Tang^{a*} and Youqing Shen^a

^aKey Laboratory of Biomass Chemical Engineering of Ministry of Education, Center for Bionanoengineering, and College of Chemical and Biological Engineering, Zhejiang University, Hangzhou, Zhejiang, 310027, China. E-mail: jianbin@zju.edu.cn; Fax: +86-571-87953993, Tel: +86-571-87951493

^bDepartment of Radiology, Sir Run Run Shaw Hospital (SRRSH) of School of Medicine, Zhejiang University, Hangzhou, Zhejiang, 310027, China.




Fig. S1. ¹H NMR spectrum (400 MHz, CDCl₃) of G1.




Fig. S2. ¹H NMR spectrum (400 MHz, CDCl₃) of G2.




Fig. S3. ^1H NMR spectrum (400 MHz, CDCl_3) of G3.




Fig. S4. ^1H NMR spectrum (400 MHz, CDCl_3) of G4.




Fig. S5. ^1H NMR spectrum (400 MHz, D_2O) of G1-Cys.




Fig. S6. ¹H NMR spectrum (400 MHz, D₂O) of G2-Cys.




Fig. S7. ¹H NMR spectrum (400 MHz, D₂O) of G3-Cys.




Fig. S8. ¹H NMR spectrum (400 MHz, D₂O) of G4-Cys.




Fig. S9. ^1H NMR spectrum (400 MHz, D_2O) of G3-OH.