

Supporting Information

Copper(II)-Catalyzed Remote Sulfonylation of Aminoquinolines with Sodium Sulfinates via Radical Coupling

Chengcai Xia,^{a*} Kai Wang,^a Jun Xu,^b Zhenjiang Wei,^a Chao Shen,^{b,d} Guiyun Duan,^a Qing Zhu^c
and Pengfei Zhang^{b*}

^a Pharmacy College, Taishan Medical University, Taian 271016, China

^b College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310036 China

^c College of Biology and Environmental Engineering, Zhejiang University of Technology, Hangzhou 310014, China


^d College of Biology and Environmental Engineering, Zhejiang Shuren University, Hangzhou 310015, China

Table of Contents


1. ¹ H and ¹³ C NMR spectra of the products...	2
--	---

1.¹H and ¹³C NMR spectra of the products


3a ¹H NMR




3a ¹³C NMR




3b ¹H NMR




3b ¹³C NMR




3c ¹H NMR




3c ¹³C NMR




3d ^1H NMR




3d ^{13}C NMR




3e ¹H NMR




3e ¹³C NMR




3f ¹H NMR




3f ¹³C NMR




3g ^1H NMR




3g ^{13}C NMR




3h ¹H NMR




3h ¹³C NMR




3i ¹H NMR




3i ¹³C NMR




3j ¹H NMR




3j ¹³C NMR




3k ¹H NMR




3k ¹³C NMR




3I ¹H NMR




3I ¹³C NMR




3m ¹H NMR




3m ¹³C NMR




3n ¹H NMR




3n ¹³C NMR




3o ¹H NMR




3o ¹³C NMR




3p ^1H NMR




3p ^{13}C NMR




3q ^1H NMR




3q ^{13}C NMR



3s ¹H NMR



3t ¹H NMR



^1H NMR

