Supplementary material

Biosynthesis, characterization and potentiality of lipopetides produced by Bacillus flexus S1 without inductive carbon sources

Chunli Wan a, Si Chen a, Lei Wen a, Xiang Liu a,*, Duu-Jong Lee b, Xue Yang c,d,*

- ^a Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, China
- ^b Department of Chemical Engineering, National Taiwan University of Science and Technology, Taipei 106, Taiwan
- ^c State Key Laboratory of Pollution Control and Resources Reuse, College of Environmental Science and Engineering, Tongji University, Shanghai 200092, China.
 - ^d Postdoctoral Research Station of Civil Engineering, Tongji University, Shanghai 200092, China
- *Corresponding author: Department of Environmental Science and Engineering, Fudan University, Shanghai, 200433, China; Tel: +86-21-65642018; Fax: +86-21-65643597; Email:

liuxiang@fudan.edu.cn (Xiang Liu); State Key Laboratory of Pollution Control and Resources Reuse, College of Environmental Science and Engineering, Tongji University, Siping Road, Shanghai 200092, China. Tel: +86-21-65984275; Fax: +86-21-65984275; E-mail: xueyang_hit@163.com (Xue Yang).



Figure S1 Colonies picture of Bacillus flexus S1.




Figure S2 Scanning electron microscope (SEM) micrograph of Bacillus flexus S1.




Figure S3 Lipopeptides produced by *Bacillus flexus* S1 fed with LB medium. Left picture is pure LB medium as control test. Right picture is LB medium cultured with *Bacillus flexus* S1, and the white precipitates in the bottom of cuvette is lipopeptides.