

Electronic Supplementary Informations

Two-stage pH-sensitive doxorubicin hydrochloride loaded core-shell nanoparticles with dual drug-loading strategies for the potential anti-tumor treatment

Xiaoyue Yu^a, Bo Zhang^a, Tianqi Wang^a, Jing Zhang^a, Shengjun Mu^a, Chunxi Liu^{*b} and Na Zhang^{*a}

^aDepartment of Pharmaceutics, School of Pharmaceutical Science, Shandong University, 44 Wenhua Road, Jinan 250012, Shandong Province, China. E-mail: zhangnancy9@sdu.edu.cn; Fax: +86 0531 88382548; Tel: +86 0531 88382015

^bPharmaceutical Department, Qilu Hospital of Shandong University, 107 Wenhua Road, Jinan 250012, Shandong Province, China. E-mail: liuchunxi1985@163.com




Fig. S1 Agarose gel electrophoresis picture with different weight ratios of PEI to CGA-ODNs from 339 : 4096 to 339 : 16 (A). Average size and PDI of POD with different weight ratios of PEI to CGA-ODNs from 339 : 256 to 339 : 16 (B).




Fig. S2 Agarose gel electrophoresis picture with different weight ratios of CMCS to CGA-ODNs from 75 : 128 to 75 : 1 (A). Zeta potentials of CPOD with different weight ratios of CMCS to CGA-ODNs from 75 : 128 to 75 : 1 (B).