

Electronic Supplementary Information

High sensitivity chlorogenic acid detection based on multiple layer-by-layer self-assembly films of chitosan and multi-walled carbon nanotubes on a glassy carbon electrode

Wenxue Cheng,^a Jianzhi Huang,^a Cheng Liu,^a Qiang Zeng,^{*a} Yanli Tong,^{*b} Lishi Wang,^a and Faliang Cheng^{*c}

a: School of Chemistry and Chemical Engineering, South China University of Technology, Guangzhou 510641, People's Republic of China.

b: Guangdong No.2 Provincial People's Hospital, Guangzhou 510317, People's Republic of China.

E-mail: ceqzeng@scut.edu.cn; tongh2008@126.com;

Tel: +86 20 87112906

*Corresponding authors.

1) Dr. Qiang Zeng, E-mail address: ceqzeng@scut.edu.cn;

2) Prof. Yanli Tong, E-mail address: tongh2008@126.com.




Fig. S1 Cyclic voltammograms of 1×10^{-4} mol dm⁻³ chlorogenic acid in phosphate buffer solution (pH 1.0) recorded on the bare electrode at different scan rates: 10, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500 and 550 mV s⁻¹ (from bottom to top). Inset: The relationship of scan rates vs. oxidation and reduction peak currents.