Synthesis and Biological Evaluation of 3-Amino-3-

hydroxymethyloxindoles as Potential Anti-cancer Agents

Kaili Jia, a Xinxin Lv, a Dong Xing, Jiuwei Che, Donglan Liu, Nilesh J. Thumar, Suzhen Dong*, Wenhao Hu*

Shanghai Engineering Research Center of Molecular Therapeutics and New Drug Development, School of Chemistry and Molecular Engineering, East China Normal University, 3663 North Zhongshan Rd., Shanghai 200062, China.

⁺ ^aThese authors contribute to this paper equally.

HPLC condition:

ACQUITY UPLC^R BEH C18 column, MeCN/H₂O = 5%_95%_4min_common; flow rate = 0.6 mL / min; [Gradient Table] Time(min) Flow Rate(mL/min) %A %B %C %D Curve 1. Initial 0.600 95.0 5.0 0.0 0.0 Initial 2. 2.80 0.600 10.0 90.0 0.0 0.0 6


3. 3.20 0.600 10.0 90.0 0.0 0.0 6

4. 3.50 0.600 100.0 0.0 0.0 0.0 6


 $5.\ 4.00\ 0.600\ 100.0\ 0.0\ 0.0\ 0.0\ 6$

Comment: A:H2O_5%ACN B:ACN t=0min,start 5%B,95%A

Flow Ramp Rate: 0.45 min



5a Sample Report:



7.094 1.924 1.004 1.874

10.5 10.0 9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0

1.00H

5. 5

2.00-J

1.00-f

5.0 4.5 4.0 3.5 3.0 2.5 2.0 fl (ppm)


1.00-I


1.5 1.0 0.5 0.0 -0.5

Sample 35 Vial 1:9 ID File LXX01-040 Date 06-Jan-2016 Time 10:50:14 Description


-200 -0


-200







5c Sample Report:





Sample 23 Vial 1:1 ID File LXX01-005 Date 05-Jan-2016 Time 22:26:11 Description





5e Sample Report:

2.048 Range: 2.053 3: UV Detector: 254 3.0 (1) 100% 2.14 2.0 AU 1.0 0.0 4.00 0.50 3.50 1.00 1.50 2.00 2.50 3.00 Peak Number Compound Time Areabs Area %Total Width Height Mass Found 1 2.14 4e+004 100.00 0 2e+006 LXX01-032 boss boss a HWH -8500 -8000 7500 -7000 -6500 -6000 5f CH3 -5500 -5000 -4500 -4000 -3500 -3000 -2500 -2000 -1500 1000 -500 0 1.04-1 0.92 0.95 1.07 € 1.03 € 3.00 ₹

9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 fl (ppn)


--500


1.05 1.02 1.01 1.04 1.05 1.05

10.5


11.5

Sample 31 Vial 1:5 ID File LXX01-029 Date 06-Jan-2016 Time 10:23:49 Description








5g Sample Report:




11.5 11.0 10.5 10.0 9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 -0.5 -1.0 fl (ppm)










5i Sample Report:




.1.0 10.5 10.0 9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 -0.5 fl (ppm)










5k Sample Report:







Sample 28 Vial 1:2 ID File LXX01-026 Date 05-Jan-2016 Time 22:32:26 Description







5m Sample Report:



Western blot

To study the underlying apoptotic mechanism, SJSA-1 cells were treated with compound **5m** at different concentration (0 μ M, 10 μ M, 20 μ M or 40 μ M) for 48 h. Then the cells were collected for western blotting analysis. The proteins of whole cell lysates were prepared in IP and western cell lysis buffer (Beyotime, China), and quantified by the bicinchoninic acid (BCA) method. Immunoblotting analysis of proteins in cell lysates was performed according to the antibody manufacturer's recommendations. Primary antibodies used were as follows: anti-Bcl-2(2872#), anti-Bax (2772#) and anti-tubulin (15115#) were purchased from Cell Signaling Technology (Danvers, MA, USA). The results showed that 5m-treatment induced an increase of anti-apoptosis protein Bcl-2 expression. And the ratio of Bax/Bcl-2 was elevated in 5a-treated cells in a dose-dependent manner, which suggested that 5a might induced cell apoptosis in SJSA-1 through an intrinsic mitochondrial pathway. Further study on the investigation of its underlying mechanism is in progress.

