Electronic Supplementary Material (ESI) for Soft Matter. This journal is © The Royal Society of Chemistry 2016

Micellization of long-chain ionic liquid in deep eutectic solvent

X. N. Tan,^{a,b} J. L. Zhang,^{*a} T. Luo,^{a,b} X. X. Sang,^a C. C. Liu,^a B. X. Zhang,^a L. Peng,^a W. Li^{*b} and B. X. Han^a

^aBeijing National Laboratory for Molecular Sciences, CAS Key Laboratory of Colloid and Interface and Thermodynamics, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

> ^bDepartment of Chemistry, Capital Normal University, Beijing 100048, China *e-mail: zhangjl@iccas.ac.cn; wli@cnu.edu.cn



Figure S1. Fluorescence spectra of pyrene in C₈mimCl/glyceline.



Figure S2. Fluorescence spectra of pyrene in C₁₀mimCl/glyceline.



Figure S3. Fluorescence spectra of pyrene in C_{12} mimCl/glyceline.



Figure S4. Fluorescence spectra of pyrene in C₁₄mimCl/glyceline.



Figure S5. Fluorescence spectra of pyrene in C_{16} mimCl/glyceline.



Figure S6. XRD pattern of the $Cu_3(BTC)_2$ synthesized in glyceline.



Figure S7. SEM (a) and TEM (b) images of the $Cu_3(BTC)_2$ synthesized in glyceline.