## Supporting Information: Strings and Stripes Formed by a Protein System Interacting via a Directional Patch Attraction

Weimin Li,<sup>†</sup> Maxim Morin,<sup>†</sup> Emil Gustavsson,<sup>†</sup> Björn A. Persson,<sup>‡</sup> Mikael Lund,<sup>‡</sup> and Malin Zackrisson Oskolkova<sup>\*,†</sup>

Division of Physical Chemistry, Lund University, POB 124, 22100 Lund, Sweden, and Division of Theoretical Chemistry, Lund University, POB 124, 22100 Lund, Sweden

E-mail: malin.zackrisson@fkem1.lu.se

<sup>\*</sup>To whom correspondence should be addressed

<sup>&</sup>lt;sup>†</sup>Division of Physical Chemistry, Lund University, POB 124, 22100 Lund, Sweden

<sup>&</sup>lt;sup>‡</sup>Division of Theoretical Chemistry, Lund University, POB 124, 22100 Lund, Sweden




Figure 1: Cryo-TEM image of lactoferrin strings at protein concentration of 80 mg/mL at ionic strength 35 mM and pH 7.




Figure 2: Cryo-TEM image of lact oferrin strings at protein concentration of 80 mg/mL at ionic strength 35 mM and pH 7.