

Overcoming multidrug resistance by a combination of chemotherapy and photothermal therapy mediated by carbon nanohorns

Junling Wang,^a Ran Wang,^a Fangrong Zhang,^b Yajun Yin,^a Leixia Mei,^a Fengjuan Song,^a Mingtao Tao^a, Wanqing Yue,^a
Wenyong Zhong^{a, c, †}


^a Department of analytical chemistry, China Pharmaceutical University, Nanjing 210009, PR China

^b State Key Laboratory of Natural Medicines, Department of Pharmaceutics, China Pharmaceutical University,
24 Tongjiaxiang, Nanjing 210009, PR China

^c Key Laboratory of Biomedical Functional Materials, China Pharmaceutical University, Nanjing 210009, PR
China

[†] Corresponding author. Tel.: +86-025-86185180; fax: +86-025-86185160

E-mail address: wyzhong@cpu.edu.cn. (W. Zhong).



$C_{29}H_{32}O_{13}$ 588.56

Fig. S1. The structure of etoposide (ETO)




Fig. S2 The histogram of A549 and A549R cells incubated with ETO@oxCNHs/PEG-PA-FITC at 37 °C for 4 h.




Fig. S3 The histogram of A549R cells incubated with ETO@oxCNHs/PEG-PA-FITC at 4 °C or 37 °C.




Fig. S4 Histologic assessments of major organs and tumor tissues with H-E staining in mice (200 \times). (A) NS; (B) oxCNHs/PEG-PA; (C) ETO; (D) ETO@oxCNHs/PEG; (E) ETO@oxCNHs/PEG-PA; (F) ETO@oxCNHs/PEG-PA + NIR.