

Engineered Highly Sensitive Whole-Cell Mercury Biosensors Based on Positive Feedback Loop from Quorum-Sensing System

Sheng Cai^{1,2}, Yifei Shen^{3,4}, Yan Zou^{1,2}, Peiqing Sun^{1,2}, Wei Wei^{1,2,*}, Jing Zhao^{1,2,*}, and Chuan Zhang^{3,4,*}

¹State Key Laboratory of Coordination Chemistry, Institute of Chemistry and BioMedical Sciences, School of Chemistry and Chemical Engineering, Collaborative Innovation Center of Chemistry for Life Sciences, Nanjing University, Nanjing, 210093, China

²State Key Laboratory of Pharmaceutical Biotechnology, School of Life Sciences, Nanjing University, Nanjing, 210093, China

³Lab of Efficient Architectures for Digital-communication and Signal-processing (LEADS), School of Information Science and Engineering, Southeast University, Nanjing, 211100, China

⁴National Mobile Communications Research Laboratory, School of Information Science and Engineering, Southeast University, Nanjing, 211100, China

*weiwei@nju.edu.cn; jingzhao@nju.edu.cn; chzhang@seu.edu.cn

Support Information

Table S1. The primers used in the article

Primers	Sequences
CmerR-1	TCTAGACTAAGGCATAGCCGAAACCTG
CPmer-2	TACTAGTGCCTGTCCCTTCGAATTG
MerR-1	ATGGAAAACAATTGGAGAA
MerR-2	AAGGCATAGCCGAAACCTGCC
Mrfp-1	ATGGTGAGCAAGGGCGAGGA
Mrfp-2	TTATTACTTGTACAGCTCGT
LuxR-1	ATGAAAAACATAATGCCGA
LuxR-2	TTATTAGTGTACACTAGCACTAT
LuxI-1	ATGACTATAATGATAAAAAAA
LuxI-2	TTATTAGAGATCTACACTAGCACTAT
Pcons2-1	TTGACGGCTAGCTCA
Pcons2-2	GCTAGCACTGTACCT
Ter-1	AAGCGACAGGCCGCCACCGGCATGGACGAGCTG
Ter-2	CGGACTAGTATAACGCAGAAAGGCCACCC