

Supplementary Information for

Catalytically Active Ceria-Supported Cobalt-Manganese Oxide Nanocatalysts for Oxidation of Carbon Monoxide

Xu Wang^{a,d}, Lin-Ying Du^b, Meng Du^{a,e}, Chao Ma^{c,}, Jie Zeng^c, Chun-Jiang Jia^{b,*}, Rui Si^{a,*}*

^a Shanghai Synchrotron Radiation Facility, Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201204, China

^b Key Laboratory for Colloid and Interface Chemistry, Key Laboratory of Special Aggregated Materials, School of Chemistry and Chemical Engineering, Shandong University, Jinan 250100, China

^c Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei, Anhui 230026, China

^d University of Chinese Academy of Science, Beijing 10049, China

^e Department of Chemistry, College of Science, Shanghai University, Shanghai 200444, China




Figure S1. TEM images of fresh (a-c) and used (d-f) ceria-supported cobalt-manganese oxide samples: (a,d) CoO_x/CeO₂; (b,e) CoMnO_x/CeO₂; (c,f) MnO_x/CeO₂.




Figure S2. HRTEM images of fresh (a-c) and used (d-f) ceria-supported cobalt-manganese oxide samples: (a,d) CoO_x/CeO₂; (b,e) CoMnO_x/CeO₂; (c,f) MnO_x/CeO₂.