

SUPPORTING INFORMATION

Vibrational energy relaxation dynamics of diatomic molecules inside superfluid helium nanodroplets. The case of the I₂ molecule

Arnau Vilà,^{a,b,‡} Miguel Paniagua,^b and Miguel González^{a,*}

^a Departament de Ciència de Materials i Química Física and IQTC, Universitat de Barcelona, c/ Martí i Franquès, 1, 08028 Barcelona (Spain).

^b Departamento de Química Física Aplicada, Universidad Autónoma de Madrid, c/ Francisco Tomás y Valiente, 7, 28049 Cantoblanco (Spain).

CONTENTS

Figure S1. Vibrational populations for the initial excitation $\nu=8$,
as a function of time. page 2

Figure S2. Vibrational populations for the initial excitation $\nu=8$,
as a function of time, considering only the $\nu=4-8$ levels. page 3

[‡] Present address: Department of Physics and Astronomy, University College London,
Gower Street, London WC1E 6BT (United Kingdom).

* Corresp. author: E-mail: miguel.gonzalez@ub.edu; Fax: +34934021231.

Figure S1. Vibrational populations for the initial excitation $v=8$, as a function of time. Please, note that the decay of this level is very fast. This can be better seen in Figure S2.




Figure S2. Vibrational populations for the initial excitation $v=8$, as a function of time, considering only the $v=4-8$ levels.

