
A Theoretical Prediction on the Shear-induced Phase Transformation of TKX-50

Zhipeng Lu,^{†,‡} Xianggui Xue,[†] and Chaoyang Zhang^{*†§}

[†] *Institute of Chemical Materials, China Academy of Engineering Physics (CAEP), P. O. Box 919-327, Mianyang, Sichuan 621900, China.*

[‡] *Department of Mathematics and Physics, Officers College of CAPF, Chengdu, 610213, China.*

[§] *Beijing Computational Science Research Center, Beijing 100048, China.*

Electronic Supplementary Information (ESI)

Table S1. Fractional Coordinates of the BCPs and Symmetry Multiplicity of TKX-50.

HB kind	D-H...A	Fraction coordinate	Symmetry Multiplicity
Intralayered	O2-H2...O1	(0.0980, 0.6366, 0.8555)	4
	N5-H4...O1	(0.5208, 0.3856, 0.1074)	4
	N5-H1...O1	(0.2279, 0.0210, 0.3905)	4
	N5-H3...N3	(0.4340, 0.3461, 0.8217)	4
	N5-H1...N4	(0.0511, 0.0790, 0.3554)	4
Interlayered	N5-H3...N2	(0.4111, 0.2539, 0.4467)	4
	N5-H3...O2	(0.2212, 0.2277, 0.2821)	4