

Self-Assemblies of TTF Derivatives Programmed by Alkyl Chain and Functional Group

Xiao-yang Zhu^{a,b†}, Bin Tu^{b†}, Guo-jun Hu^c, Qiao-jun Fang^b, Jun-jie Qi^{a*}, Xun-wen Xiao^{c*}, Yan-fang
Geng^{b*} and Qing-dao Zeng^{b*}

^a School of Materials Science and Engineering, University of Science and Technology Beijing, Beijing 100083, China. E-mail: junjieqi@ustb.edu.cn

^b CAS Key Laboratory of Standardization and Measurement for Nanotechnology, CAS Center for Excellence in Nanoscience, National Center for Nanoscience and Technology (NCNST), 11 Zhongguancun Beiyitiao, Beijing 100190, China. E-mail: zengqd@nanoctr.cn, gengyf@nanoctr.cn

^c College of Chemical Engineering, Ningbo University of Technology, Ningbo 315211, P. R. China. E-mail: xunwenxiao@nbut.edu.cn

[†] These authors contributed equally to this work.




Figure S1. (a) Observed STM image of HOPG at a scanning condition ($I_{\text{set}} = 600$ pA, $V_{\text{bias}} = 10$ mV). (b) Illustration of the graphite lattice.