

SUPPORTING INFORMATION

THERMO-ENHANCED RING-OPENING POLYMERIZATION OF ϵ -CAPROLACTONE: THE SYNTHESIS, CHARACTERIZATION AND CATALYTIC BEHAVIOR OF ALUMINUM HYDROQUINOLIN-8-OLATES

Qiurui Zhang,^{a,b,c} Wenjuan Zhang,^{*,a} Natesan Mannangatti Rajendran,^b Tongling Liang,^b and Wen-Hua Sun^{*,b,c}

^aSchool of Materials Science and Engineering, Beijing Institute of Fashion Technology, Beijing 100029, China

^bKey Laboratory of Engineering Plastics, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

^c CAS Research/Education Center for Excellence in Molecular Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

Corresponding author: W.-H.Sun, whsun@iccas.ac.cn; W. Zhang, zhangwj@bift.edu.cn

Figure S1. The ¹H NMR and ¹³C NMR spectrum of **L1**

Figure S2. The ¹H NMR and ¹³C NMR spectrum of **L2**

Figure S3. The ¹H NMR and ¹³C NMR spectrum of **L3**

Figure S4. The ¹H NMR and ¹³C NMR spectrum of **L4**

Figure S5. The ¹H NMR and ¹³C NMR spectrum of **L5**

Figure S6. The ¹H NMR and ¹³C NMR spectrum of **C1**

Figure S7. The ¹H NMR and ¹³C NMR spectrum of **C2**

Figure S8. The ¹H NMR and ¹³C NMR spectrum of **C3**

Figure S9. The ¹H NMR and ¹³C NMR spectrum of **C4**

Figure S10. The ¹H NMR and ¹³C NMR spectrum of **C5**

Figure S11. The ¹H NMR and ¹³C NMR spectrum of **C6**

Figure S12. The ¹H NMR and ¹³C NMR spectrum of **C7**

Figure S13. The ¹H NMR and ¹³C NMR spectrum of **C8**

Figure S14. The ¹H NMR and MALDI-TOF spectrum of PCL by **C3**

Figure S15. The ¹H NMR and MALDI-TOF spectrum of PCL by **C4**

Figure S16. The ¹H NMR and MALDI-TOF spectrum of PCL by **C7**




Figure S1. The ¹H NMR and ¹³C NMR spectrum of L1





Figure S2. The ^1H NMR and ^{13}C NMR spectrum of L2




Figure S3. The ^1H NMR and ^{13}C NMR spectrum of L3





Figure S4. The ¹H NMR and ¹³C NMR spectrum of L4





Figure S5. The ^1H NMR and ^{13}C NMR spectrum of L5




Figure S6. The ^1H NMR and ^{13}C NMR spectrum of C1





Figure S7. The ¹H NMR and ¹³C NMR spectrum of C2





Figure S8. The ^1H NMR and ^{13}C NMR spectrum of C3





Figure S9. The ¹H NMR and ¹³C NMR spectrum of C4





Figure S10. The ¹H NMR and ¹³C NMR spectrum of C5




Figure S11. The ^1H NMR and ^{13}C NMR spectrum of C6




Figure S12. The ^1H NMR and ^{13}C NMR spectrum of C7





Figure S13. The ¹H NMR and ¹³C NMR spectrum of **C8**





Figure S14. The ¹H NMR and MALDI-TOF spectrum of PCL by C3





Figure S15. The ^1H NMR and MALDI-TOF spectrum of PCL by C4





Figure S16. The ^1H NMR and MALDI-TOF spectrum of PCL by C7