

Table 1 Supplemental: Identified extractives isolated upon successive extractions with dichloromethane (D), toluene/ethanol (T-E) and acetone (A). Solvents in which the identified extractives were found are indicated.

Identified Extractive ^{a,b}	Amount [%]	Fraction
Hexadecanoic acid	1.6	D, T-E, A
1,3-Dimethoxy-5-[(1E)-2-phenylethenyl] benzene	2.0	D, T-E
3-Hydroxy-4-methoxybenzyl alcohol	2.3	D
Pimaric acid	9.0	D, A
Isopimaric acid	10.5	D, T-E
Dehydroabietic acid	9.5	D, A
Abietic acid	10.7	D, T-E, A
5- α -Pregnan-3- β ,16- α ,20- α -triol	0.7	D, T-E, A
5-Ethyl-5-isopropyl-2,4,6(1H,3H,5H)-pyrimidinetrione	0.9	D
Dehydroabietic acid	1.7	D, T-E, A
Methyl abieta-8,11,13-trien-18-oate	2.3	D
Silane, diethylpentadecyloxy(3-phenylpropoxy)-	0.9	D
4',5,7-trihydroxyflavanone	0.8	D
3,21-Dihydroxy-3- β -pregn-5-en-20-one	2.6	D, T-E, A
17- α -3-Hydroxypregnenolone	0.8	D
7-Oxodehydroabietic acid	3.2	D, T-E
21-Hydroxyprogesterone	0.5	D
Docosa-8,14-diyne- <i>cis</i> -1,22-diol	0.6	D
Propanoic acid	0.6	T-E
4-Hydroxy-3-methoxybenzaldehyde	0.7	A
Benzaldehyde	0.7	T-E
Benzylmalonic acid	2.1	T-E
3,5-Dimethyl-2-cyclohexen-1-one	2.8	A
(R)-3-Pyrrolidinol	0.5	A
3,5-Dihydroxybenzoic acid	0.8	T-E, A
5-Methyl-4-nonene	1.9	A
2,6-Dihydroxybenzoic acid	2.0	T-E
4-Hydroxy-3-methoxybenzyl alcohol	5.3	T-E, A
4'-Methoxy-2-hydroxystilbene	1.6	T-E
1-Heptadecene	0.5	T-E
4-Hydroxy-3-methoxybenzoic acid	4.9	T-E, A
1-Dodecene	0.6	T-E
5-Methyl-4-undecene	1.0	A
1-Pentadecene	0.7	T-E
α -Pinene	0.5	T-E
Heneicosyl acetate	0.8	A
Azelaic acid	0.6	A
1-Dodecanol	0.7	A
2-Deoxy ribose	1.4	A
4-Hydroxy-benzenepropionic acid	0.5	A
2-Methyl-2(<i>p</i> -methoxy)mandelate	1.0	A
5-Dehydro- <i>epi</i> -androsterone isocaproate	0.8	A
2,2-Dihydroxy-(4-hydroxyphenyl)propane	0.9	A
5-Allyl-1-methoxy-2,3-dihydroxybenzene	1.2	A
3,7-Dimethyl-9-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2E,4E,6E,8E-nonatetraenoic acid	0.7	A
4-Hydroxy-3-methoxycinnamaldehyde	0.7	A
Methyl 4-hydroxycinnamate	3.7	A
Oleic acid	2.7	D, A
β -Sitosterol	3.8	D
(E)-9-Octadecenoic acid	1.7	T-E
(Z,Z)-9,12-Octadecadienoic acid	2.0	D, T-E, A
(-)-Nortrachelogenin	1.4	T-E

Naringenin	0.9	T-E
Octadecanoic acid	1.9	D, T-E, A
Docosanoic acid	1.3	D
Tetracosan-1-ol	2.0	D

a: Analyses was carried out after silylation of the free hydroxyl groups with *N,O*-bis(trimethylsilyl)trifluoroacetamide.

b: Extractives with a relative abundance of >0.5% are listed.