

Electronic Supplementary Information

Low energy recycling of ionic liquids via freeze crystallization during cellulose spinning

Yanrong Liu,^{a,b,c} Anne S. Meyer,^c Yi Nie,^a Suojiang Zhang,^{*a} Kaj Thomsen^{*b}

^aBeijing Key Laboratory of Ionic Liquids Clean Process, Key Laboratory of Green Process and Engineering, State Key Laboratory of Multiphase Complex Systems, Institute of Process Engineering, Chinese Academy of Sciences, P. O. Box 353, Beijing, 100190, China, E-mail: sjzhang@ipe.ac.cn.

^bCenter for Energy Resources Engineering (CERE), Department of Chemical and Biochemical Engineering, Technical University of Denmark, Søltofts Plads 229, 2800, Lyngby, Denmark, E-mail: kth@kt.dtu.dk.

^cCenter for Bioprocess Engineering, Department of Chemical and Biochemical Engineering, Technical University of Denmark, Søltofts Plads 227, 2800, Lyngby, Denmark, E-mail: am@kt.dtu.dk.

1. EmimAc and EmimDep recycling processes by evaporation method




Figure 6 EmimAc recycling procedure in cellulose spinning process. H₁-H₈ are the enthalpies of the

mixtures described in each box (a) to box (h).




Figure 7 EmimDep recycling procedure in cellulose spinning process. H_1 - H_8 are the enthalpies of the mixtures described in each box (a) to box (h).