

Electronic Supplementary Information

Platinum(II)-chloroquine complexes are antimarial agents against blood and liver stages by impairing mitochondrial function

Taís S. Macedo,^{a‡} Wilmer Villarreal,^{b‡} Camila C. Couto,^a Diogo R. M. Moreira,^a Maribel Navarro,^c

Marta Machado,^d Miguel Prudêncio,^d Alzir A. Batista,^b Milena B. P. Soares^{a,e*}

^aFIOCRUZ, Instituto Gonçalo Moniz, 40296-710, Salvador, BA, Brazil.

^bUFSCAR, Departamento de Química, 13565-905, São Carlos, SP, Brazil.

^cUFJF, ICE, Departamento de Química, 36036-900, Juiz de Fora, MG, Brazil.

^dInstituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisboa, Portugal.

^eCentro de Biotecnologia e Terapia Celular, Hospital São Rafael, 41253-190, Salvador, BA, Brazil.

*Corresponding author: FIOCRUZ, Instituto Gonçalo Moniz, CEP 40296-710, Salvador, BA, Brazil.
(+55)71-31762292; email: milena@bahia.fiocruz.br

‡Footnote: Taís S. Macedo and Wilmer Villarreal share first authorship.







Figure S1: $^{31}\text{P}\{\text{H}\}$ spectra of platinum complexes in a mixture DMSO:Trizma 60:40 (D_2O). Incubation times are indicated in the right axis of each spectrum.





Figure S2: Hemolysis in uninfected erythrocytes after 1 h incubation of compounds at indicated concentration. DMSO at 0.5 % (v/v) in PBS, Saponin at 1 % v/v in PBS. Two independent experiments were performed. Error bars represent S.D. from pooling triplicate values of one experiment. *** $p < 0.01$ (one-way ANOVA and Newman-Keuls test).

Table S1: Cytostatic activity of platinum complexes against intraerythrocytic *P. falciparum*.

Comp.	<i>P. falciparum</i> , IC ₅₀ ±S.D.(μM) ^a	
	CQ-sensitive 3D7	CQ-resistant W2
WV-48	> 5.0	3.6±0.3
WV-31	> 5.0	N.D.
WV-51	>5.0	N.D.
WV-50	3.9±0.5	2.4±0.3

^a Determined 72 h after incubation with compounds.




A**B****C**

Figure S3: Ultrastructural effects of platinum complex and chloroquine in *P. berghei*-infected mice. Representative transmission electron micrographs of untreated (panel A), chloroquine at 66 µmol/kg (panel B) and WV-90 at 66 µmol/kg (panel C) after 6 h treatment. Abbreviation: N, nucleus; M, mitochondria; DV, digestive vacuole; AP, autophagosome; MP, parasitophorous vacuole. Arrows indicate hemozoin.