

Electronic Supplementary Information

A Ni-2,2'-bisdipyrinato complex as a potential sensitizer: synthesis and photoelectrochemical characterization

Alessandro Poma,^a Ivan Grigioni,^a Maria Vittoria Dozzi,^a Stéphane A. Baudron,^b Lucia Carlucci,^a
Mir Wais Hosseini^b and Elena Selli^{a*}

- ^{a.} Dipartimento di Chimica, Università degli Studi di Milano, Via Golgi 19, I-20133 Milano, Italy, elena.selli@unimi.it.
- ^{b.} Laboratoire de Tectonique Moléculaire, UMR CNRS 7140, icFRC, Université de Strasbourg, F-67000, Strasbourg.

Figure S1 – ¹ H NMR spectrum of complex Ni-bisdpmCOOH.....	2
Figure S2 – Effect of the scan rate on the first oxidation peaks of NiTCPP.....	2
Figure S3 – UV-Vis-NIR absorption spectrum of TiO ₂ deposited on FTO.....	3
Figure S4 – Difference absorption spectrum of NiTCPP adsorbed on TiO ₂	3




Figure S1 – ^1H NMR spectrum of complex Ni-bisdpmCOOH.




Figure S2 – Effect of the scan rate on the first oxidation peaks of NiTCPP.




Figure S3 – UV-Vis-NIR absorption spectrum of TiO₂ deposited on FTO.




Figure S4 – Difference absorption spectrum of NiTCPP adsorbed on TiO₂.