

Efficient synthesis and characterization of novel indolizines: exploration of *in vitro* COX-2 inhibitory activity and molecular modelling studies

Sandeep Chandrashekarappa,^a Katharigatta N. Venugopala,^{*bc} Christophe Tratat,^c Fawzi M. Mahomoodally,^d Bandar E Aldhubiab,^c Michelyne Haroun,^c Rashmi Venugopala,^e Mahendra K. Mohan,^a Rashmi S. Kulkarni,^f Mahesh V. Attimarad,^e Sree Harsha,^c Bharti Odhav,^b

Submitted to

New Journal of Chemistry (Manuscript Reference NJ-ART-12-2017-005010)

Compounds code	Description	Page number
2a	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 7-cyano-2-ethyl-3-(4-fluorobenzoyl)indolizine-1-carboxylate	4-5
2b	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 3-(4-bromobenzoyl)-7-cyano-2-ethylindolizine-1-carboxylate	6-7
2c	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Diethyl 3-(4-chlorobenzoyl)-7-cyanoindolizine-1,2-dicarboxylate	8-9
2d	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 7-cyano-3-(4-cyanobenzoyl)-2-methylindolizine-1-carboxylate	10-11
2e	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 3-(4-bromobenzoyl)-7-cyanoindolizine-1-carboxylate	12-13
2f	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 7-cyano-3-(4-methoxybenzoyl)-2-methylindolizine-1-carboxylate	14-15
2g	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 3-(4-bromobenzoyl)-7-cyano-2-methylindolizine-1-carboxylate	16-17
2h	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 7-cyano-2-ethyl-3-(4-methoxybenzoyl)indolizine-1-carboxylate	18-19
2i	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 3-(4-chlorobenzoyl)-7-cyano-2-ethylindolizine-1-carboxylate	20-21
2j	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Diethyl 7-cyano-3-(4-cyanobenzoyl)indolizine-1,2-dicarboxylate	22-23
2k	FT-IR, NMR (^1H & ^{13}C) and LC-MS of Ethyl 7-cyano-3-(4-cyanobenzoyl)indolizine-1-carboxylate	24-25











































