

Electronic Supplementary Information

Carrier-free, Self-assembled Pure Drug Nanorods Composed of 10-Hydroxycamptothecin and Chlorin e6 for Combinatorial Chemo-Photodynamic Antitumor Therapy *in vivo*

Table of Contents

Figure S1. TEM image of (A) pure HCPT nanostructure prepared through reprecipitation method and (B) self-assembled nanoparticles of deprotonated Ce6 in aqueous solution.

Figure S2. Representative hydrodynamic diameters of HCPT/Ce6 nanostructures with different ratios determined by DLS.

Figure S3. Zeta potential of HCPT/Ce6 NRs.

Figure S4. Fluorescence emission spectra of SOSG solution in the presence of HCPT/Ce6 NRs with the increase of laser irradiation time.

Figure S5. Cell viability curves and the IC_{50} values (μM) of (A) 4T1, (B) A549 and (C) MCF-7 cells.

Figure S6. Viability of (A) A549 cells (B) MCF-7 cells.

Figure S7. Representative digital photos of (A) full-scale view of mice in each group, (B) enlarged local view of tumor sites from (A), (C) excised 4T1 tumor tissues after various treatments for 12 days.




Figure S1. TEM image of (A) pure HCPT nanostructure prepared through reprecipitation method and (B) self-assembled nanoparticles of deprotonated Ce6 in aqueous solution.




Figure S2. Representative hydrodynamic diameters of HCPT/Ce6 nanostructures with different ratios determined by DLS.




Figure S3. Zeta potential of HCPT/Ce6 NRs.




Figure S4. Fluorescence emission spectra of SOSG solution in the presence of HCPT/Ce6 NRs with the increase of laser irradiation time.




Figure S5. Cell viability curves and the IC₅₀ values (μM) of (A) 4T1, (B) A549 and (C) MCF-7 cells after incubation with free HCPT or HCPT/Ce6 NRs for 48 h.




Figure S6. Viability of (A) A549 cells (B) MCF-7 cells after incubation with various formulations with (+Laser) or without laser irradiation ($\lambda = 655 \text{ nm}$, 150 mW/cm^2 , 2 min).




Figure S7. Representative digital photos of (A) full-scale view of mice in each group, (B) enlarged local view of tumor sites from (A), (C) excised 4T1 tumor tissues after various treatments for 12 days.