


Ir-catalyzed asymmetric hydrogenation of β -keto esters with chiral ferrocenyl P,N,N-ligands

Xiu-Shuai Chen, Chuan-Jin Hou,* Chao Qin, Hongzhu Liu, Yan-Jun Liu, De-Zhi Huang and Xiang-Ping Hu*


*School of Light Industry and Chemical Engineering, Dalian Polytechnic University, Dalian 116034, China. E-mail:
houcj@dlpu.edu.cn*

Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China. E-mail: xiangping@dicp.ac.cn


(S)-methyl 3-hydroxy-3-phenylpropanoate 3a




(S)-methyl 3-(2-chlorophenyl)-3-hydroxypropanoate 3b




(S)-methyl 3-(3-chlorophenyl)-3-hydroxypropanoate 3c




(S)-methyl 3-(4-chlorophenyl)-3-hydroxypropanoate 3d




(S)-methyl 3-(4-bromophenyl)-3-hydroxypropanoate 3e




(S)-methyl 3-(4-fluorophenyl)-3-hydroxypropanoate 3f




(S)-methyl 3-hydroxy-3-(*p*-tolyl)propanoate 3g




(S)-methyl 3-(4-methoxyphenyl)-3-hydroxypropanoate 3h




(S)-methyl 3-hydroxy-3-(naphthalen-2-yl)propanoate 3i




(S)-methyl 3-(furan-2-yl)-3-hydroxypropanoate 3j




(S)-methyl 3-hydroxy-3-(thiophen-2-yl)propanoate 3k



(S)-methyl 3-hydroxy-4-methylpentanoate 3l



(S)-methyl 3-hydroxybutanoate 3m






7.3873
7.3716
7.3549
7.3351
7.3087
7.3035
7.2956
7.2873
7.2574


5.1549
5.1449
5.1328
5.1229


-3.7242
2.8076
2.7855
2.7667
2.7445
2.7384
2.7283
2.6974
2.6874


HCJ-2387
PROTON CDCl₃ {D:\NMR400\02T2} nmr 48































HCJ-2396
PROTON CDC13 {D:\NMR400\02T2} nmr 57



HCJ-20170216

PROTON CDC13 {D:\NMR400\02T2} nmr 35

