

Supplementary information for:

**Surface Spermidine Functionalized PEGylated Poly (lactide-co-glycolide)
Nanoparticles for Tumor-targeted Drug Delivery**

Jianming Li^a, Juan Mao^a, Jing Tang^b, Guo Li^a, Fengling Fang^a, Yana Tang^a,
Jinsong Ding^{a, *}

^a Xiangya School of Pharmaceutical Sciences, Central South University, Changsha 410013, Hunan, China.

^b Department of Pharmaceutics, Changsha Medical University, Changsha 410219, Hunan, China




Fig.S1 (A) Scheme of the reaction between fluorescamine and a primary amine; (B) Fluorescence spectra of Boc-SPD calibrating solutions; (C) Standard curve of fluorescence intensity to Boc-SPD concentration at 480 nm.

* Address for correspondence: Xiangya School of Pharmaceutical Sciences, Central South University, Changsha 410013, Hunan, China.
E-mail addresses: dingjs0221@163.com; Fax: +86-731-82650442; Tel: +86-731-82650250




Fig.S2 *In vitro* release profiles of DOX from DOX-NPs and SPD-DOX-NPs with different SPD contents on NPs surface in PBS (pH 7.4) with 0.05 % SDS at 37 °C for 192 h. Data were shown as mean \pm SD (n=3).




Fig.S3 *In vivo* cytotoxicity of DFMO on A549 cells for 24 h. Data were shown as mean \pm SD (n=4)




Fig.S4 *In vivo* cytotoxicity of blank NPs and SPD-NPs on A549 cells according to polymer concentrations for 24 h. Data were shown as mean \pm SD (n=4)