

Supplementary information for the manuscript:

Tyrosinase mediated oxidative functionalization in the synthesis of DOPA peptidomimetics with anti-Parkinson activity

Bruno M. Bizzarri,^a Alessandro Martini,^b Francesco Serafini,^a Daniela Aversa,^b Davide Piccinino,^a Lorenzo Botta,^a Nicola Berretta,^b Ezia Guatteo,^{b,c} Raffaele Saladino^{a,*}

^a Department of Ecological and Biological Sciences (DEB), University of Tuscia, Via S Camillo de Lellis, 01100 Viterbo, Italy. saladino@unitus.it.



^b Laboratory of Experimental Neurology, Fondazione Santa Lucia IRCCS, Via del Fosso di Fiorano 64, 00143 Roma, Italy.

^c University of Naples Parthenope, Via Medina 40, 80133 Napoli, Italy



This file includes:

SI#1: ¹H NMR and ¹³C NMR spectra of DOPA peptidomimetics **2-9**



OMe-Gly-N-Boc-DOPA-OMe (**2**)





N-Boc-Gly-N-Boc-Dopa-Ome (**3**)






OMe-Val-N-Boc-Dopa-OMe (**4**)






N-Boc-Val-N-Boc-DOPA-OMe(5)






Gly-N-C-Dopa (6)



Val-N-C-Dopa (7)



Gly-O-C-Dopa (8)



Val-O-C-Dopa (9)

