


## Support Information

### A designable aminophenylboronic acid functionalized magnetic Fe<sub>3</sub>O<sub>4</sub>/ZIF-8/APBA for specific recognition of glycoproteins and glycopeptides

Shanshan Li, Dongyan Li\*, Long Sun, Yuewei Yao and Cheng Yao\*

School of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816, China.

E-mail: lidongyan111@njtech.edu.cn; [yaocheng@njtech.edu.cn](mailto:yaocheng@njtech.edu.cn)



**FigureS1.** FT-IR spectra of  $\text{Fe}_3\text{O}_4$  (a),  $\text{Fe}_3\text{O}_4/\text{ZIF-8}$  (b), and  $\text{Fe}_3\text{O}_4/\text{ZIF-8/APBA NPs}$  (c).




Figure S2.  $\text{N}_2$  sorption–desorption isotherms of the as-prepared  $\text{Fe}_3\text{O}_4/\text{ZIF-8}$  ( a ) and  $\text{Fe}_3\text{O}_4/\text{ZIF-8/APBA}$  ( b ) NPs measured at 77 K.

Table S1. Detailed information of the glycopeptides enriched by Fe<sub>3</sub>O<sub>4</sub>/ZIF-8/APBA from tryptic digestion of HRP.

Peak number	Observed m/z	Glycan composition	Amino acid sequence <sup>[a]</sup>
H1	2533	FucGlcNAc	SFAN#STQTFFNAFVEAMDR
H2	2594	XylMan3FucGlcNAc2	PTLN#TTYLQTLR
H3	2851	FucGlcNAc	GLIQSDQELFSSPN#ATDTIPLV R
H4	3323	XylMan3FucGlcNAc2	QLTPTFYDN SCPN#VSNIVR
H5	3353	XylMan3FucGlcNAc2	SFAN#STQTFFNAFVEAMDR
H6	3671	XylMan3FucGlcNAc2	GLIQSDQELFSSPN#ATDTIPLV R

[a] The *N*-glycosylation sites are marked with N#. GlcNAc = *N*-acetylglucosamine, Man = mannose, Gal =galactose.