Supporting Information For

High Cetane Renewable Diesel Fuels Prepared from Methyl Ketones and Diols

Kale W. Harrison and Benjamin G. Harvey*

US NAVY, NAWCWD, Research Department, Chemistry Division, China Lake, California 93555

Contents

Figure S1.	¹ H NMR spectrum of 2,4,5-trimethyl-2-undecyl-1,3-dioxolane (1)	S2
Figure S2.	¹³ C NMR spectrum of 2,4,5-trimethyl-2-undecyl-1,3-dioxolane (1)	S2
Figure S3.	¹ H NMR spectrum of 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)	S3
Figure S4.	¹³ C NMR spectrum of 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)	S3
Figure S5.	¹ H NMR spectrum of 2-methyl-2-undecyl-1,3-dioxolane (3)	S4
Figure S6.	¹³ C NMR spectrum of 2-methyl-2-undecyl-1,3-dioxolane (3)	S4
Figure S7.	DSC data for 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)	S5
Figure S8.	DSC data for 2-methyl-2-undecyl-1,3-dioxolane (3)	S5



Figure S1. ¹H NMR spectrum of 2,4,5-trimethyl-2-undecyl-1,3-dioxolane (1)



Figure S2. ¹³C NMR spectrum of 2,4,5-trimethyl-2-undecyl-1,3-dioxolane (1)



Figure S3. ¹H NMR spectrum of 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)



Figure S4. ¹³C NMR spectrum of 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)



Figure S5. ¹H NMR spectrum of 2-methyl-2-undecyl-1,3-dioxolane (3)



Figure S6. ¹³C NMR spectrum of 2-methyl-2-undecyl-1,3-dioxolane (**3**)



Figure S7. DSC data for 2,4-dimethyl-2-undecyl-1,3-dioxolane (2)



Figure S8. DSC data for 2-methyl-2-undecyl-1,3-dioxolane (3)