

Supporting Information

Enhanced hydrogen production from ammonia borane using controlled plasmonic performance of Au nanoparticles deposited on TiO₂

Seongwon Jo,^a Priyanka Verma,^b Yasutaka Kuwahara,^{bc} Kohsuke Mori,^{bcd}
Wonyong Choi^{*a} and Hiromi Yamashita^{*bc}

^aDivision of Environmental Science and Engineering, Dept. of Chemical Engineering,
Pohang University of Science and Technology (POSTECH), Pohang 37673, Korea

^bDivision of Materials and Manufacturing Science, Graduate School of Engineering,
Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

^cElements Strategy Initiative for Catalysts & Batteries Kyoto University (ESICB), Kyoto
University, Katsura, Kyoto 615-8520, Japan


^dJST, PRESTO, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012, Japan

*To whom correspondence should be addressed.

E-mail: wchoi@postech.edu; Tel.: +82-54-279-2283; Fax: +82-54-279-8299

E-mail: yamashita@mat.eng.osaka-u.ac.jp; Tel.: +81-6-6879-7457; Fax: +81-6-6879-7

(a)



(b)




Fig. S1. XPS (a) Ti 2p and (b) O 1s spectra of TiO₂ and Au1/TiO₂.