## **Electronic Supplementary Information (ESI)**

Oral delivery of curcumin *via* porous polymeric nanoparticles for effective ulcerative colitis therapy

Qiubing Chen,<sup>‡a</sup> Xiaoying Si,<sup>‡a</sup> Lijun Ma,<sup>a</sup> Panpan Ma,<sup>a</sup> Meili Hou,<sup>a</sup> Shuang Bai,<sup>a</sup> Xiaoshuai Wu,<sup>a</sup> Ying Wan<sup>\*b</sup>, Bo Xiao<sup>\*a,c</sup> and Didier Merlin<sup>c,d</sup>

<sup>a</sup>Institute for Clean Energy and Advanced Materials, Faculty of Materials and Energy, Southwest University, Chongqing, 400715, P. R. China
<sup>b</sup>College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, P. R. China
<sup>c</sup>Center for Diagnostics and Therapeutics, Institute for Biomedical Science, Georgia State University, Atlanta, 30302, USA

<sup>d</sup>Atlanta Veterans Affairs Medical Center, Decatur, 30033, USA


<sup>‡</sup>These authors contributed equally to this work.

\*Author for correspondence

Dr. Ying Wan, Email: ying\_wan@hust.edu.cn

Dr. Bo Xiao, Email: bxiao@gsu.edu

Supplementary Figure 1. *In vitro* cytotoxicity of non-porous NPs and porous NPs against Raw 264.7 macrophages (a) and CT-26 cells (b) after co-incubation for 24 h based on MTT assays. Each point represents the mean  $\pm$  S.E.M. (n=5).



## Supplementary Fig. 1

Chen Q. et al.