

Supplementary Information

Inflammatory activation of human serum albumin- or ovalbumin-modified chitosan particles to macrophages and their immune response in human whole blood

Yixian Zhang^a, Wang Du^a, Kathrin Smuda^b, Radostina Georgieva^b, Hans Bäumler^{b*}, Changyou Gao^{a,c*}

^a *MOE Key Laboratory of Macromolecular Synthesis and Functionalization, Department of Polymer Science and Engineering, Zhejiang University, Hangzhou 310027, China*

^b *Institute of Transfusion Medicine and Berlin-Brandenburg Center for Regenerative Therapies, Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany*

^c *Dr. Li Dak Sum & Yip Yio Chin Center for Stem Cell and Regenerative Medicine, Zhejiang University, Hangzhou 310058, China*

*Corresponding authors:

Email: cygao@mail.hz.zj.cn (C.Y. Gao)

E-mail: hans.baeumler@charite.de (H. Bäumler)

Fluorescence Spectra of proteins-modified CS particles




Figure S1 Fluorescence spectra of different CS particles measured by fluorescence spectroscopy (LS55, Perkin Elmer, Japan) with the excitation wavelength of 488 nm. The particles were dispersed in water to a concentration of 100 $\mu\text{g/mL}$ with the assistance of a bath sonicator.