Supporting Information

Black TiO₂ based nanoprobes for T₁-weighted MRI guided photothermal

therapy in CD133 high expressed pancreatic cancer stem-like cells

Siqi Wang^{†a}, Wenzhi Ren^{†b,c}, Jianhua Wang^{*a}, Zhenqi Jiang^{b,c}, Madiha Saeed^{b,c}, Lili

Zhang^d, Aiguo Li^d, Aiguo Wu^{*b}

^aDepartment of Radiology, The Affiliated Hospital of Medical School of Ningbo University, Ningbo University School of Medicine, Ningbo, 315020, Zhejiang Province, China.

^bCAS Key Laboratory of Magnetic Materials and Devices & Key Laboratory of Additive Manufacturing Materials of Zhejiang Province& Division of Functional Materials and Nanodevices, Ningbo Institute of Materials Technology and Engineering, Chinese Academy of Sciences, 1219 ZhongGuan West Road, 315201, Ningbo, China

^cUniversity of Chinese Academy of Sciences, No. 19(A) Yuquan Road, Shijingshan District, Beijing, 100049, P.R. China

^dShanghai Synchrotron Radiation Facility, Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai, 201204, China

*Corresponding Author. Jianhua Wang E-mail address: woxingw@sina.com

*Corresponding Author. Aiguo Wu E-mail address: aiguo@nimte.ac.cn



Figure S1. 1H NMR spectra were obtained for quantitative analysis of NH₂-PEG-COOH. Sodium acetate as internal standard (δ 1.817) substances, δ 3.610 belongs to NH₂-PEG-COOH, and D₂O as solvent.