

Supporting Information

Dynamics of Benzene Excimer Formation from the Parallel-Displaced Dimer

Thiago Messias Cardozo,^{1,*} Andre Pessoa Galliez,¹ Itamar Borges Jr.,² Felix Plasser,³ Adelia J. A. Aquino,^{4,5,6} Mario Barbatti,^{7,*} Hans Lischka,^{4,5,6,*}

¹ Instituto de Química, Universidade Federal do Rio de Janeiro,
Rio de Janeiro, RJ 21941-909, Brazil.

² Departamento de Química, Instituto Militar de Engenharia,
Praça General Tibúrcio, 80, 22290-270 Rio de Janeiro, Brazil

³ Department of Chemistry, Loughborough University, LE11 3TU, United Kingdom

⁴ School of Pharmaceutical Sciences and Technology, Tianjin University,
Tianjin 300072, P.R. China

⁵ Department of Chemistry and Biochemistry, Texas Tech University,
Lubbock, Texas 79409, United States

⁶ Institute of Theoretical Chemistry, University of Vienna,
Währingerstraße 17, 1090 Vienna, Austria

⁷ Aix Marseille Univ, CNRS, ICR, Marseille, France.

* Corresponding authors. E-mail addresses: TMC: thiago@iq.ufrj.br; MB:
mario.barbatti@univ-amu.fr; HL: hans.lischka@univie.ac.at




Figure S1. Average total kinetic energy of hydrogen atoms during the dynamics. The values are divided by the $T(\text{Hyd})$ at 0fs.