Supplementary materials for "Troxerutin Subdues Hepatic Tumorigenesis Disrupting MDM2-p53 Interaction

Nisha Susan Thomas^{a*}, Kiran George^b, Athavan Alias Anand Selvam^c

^a Department of Biochemistry and Biotechnology, Annamalai University, Tamil Nadu, India.

^b Department of Electronics and Instrumentation Engineering, Bioengineering, Annamalai University, Tamil Nadu, India

^c Department of Organic Chemistry, Indian Institute of Science, Bangalore, India.

Supplementary fig. 1: Quantitative deconvoluted images (analyzed in ImageJ software) of immunohistochemical expressions of GST-P foci and PCNA positive cells showing the intensity of DAB in the liver sections of control and experimental rats.



Supplementary fig. 2: Quantitative deconvoluted images (analyzed in ImageJ software) of immunohistochemical expressions of PARP-1 showing the intensity of DAB in the liver sections of control and experimental rats.



Supplementary fig. 3: Quantitative deconvoluted images (analyzed in ImageJ software) of immunohistochemical expressions of MDM2, p53, Bcl-2 and Bax showing the intensity of DAB in the liver sections of control and experimental rats.



Supplementary fig. 4: Quantitative deconvoluted images (analyzed in ImageJ software) of immunohistochemical expressions of cytochrome C, caspase-9 and caspase-3 showing the intensity of DAB in the liver sections of control and experimental rats.

