

New Journal of Chemistry
Electronic Supplementary Information
to

**Cd^{II}/Zn^{II} discrimination by 2,5-diphenyl[1,3,4]oxadiazole based
fluorescent chemosensors**

Mauro Formica,^a Gianluca Ambrosi,^a Vieri Fusi,^{a*} Luca Giorgi,^{a*} Massimiliano Arca,^b Alessandra Garau,^b Anna Pintus,^b Vito Lippolis^b

^aDepartment of Pure and Applied Sciences, University of Urbino, P.zza Rinascimento 6, I-61029 Urbino, Italy. Tel/Fax: +39-0722-304884. Email: luca.giorgi@uniurb.it

^bDipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Cagliari, S.S. 554 Bivio per Sestu, 09042 Monserrato (CA), Italy




Figure S1. ^1H -NMR titration of **L1** in CD_3CN at 298.1 K obtained by adding several amounts of $\text{Zn}(\text{ClO}_4)_2$ to a **L1** ($1.0 \cdot 10^{-2} \text{ mol} \cdot \text{dm}^{-3}$) solution.




Figure S2. ¹H-NMR titration of **L1** in in CD₃CN at 298.1 K obtained by adding several amounts of Cd(ClO₄)₂ to a **L1** ($1.0 \cdot 10^{-2} \text{ mol} \cdot \text{dm}^{-3}$) solution.




Figure S3. Selected KS-orbital drawings calculated for **L1** at the DFT level. Contour value= 0.05 e.




Figure S4. Selected KS-orbital drawings calculated for **L2** at the DFT level. Contour value= 0.05 e.